



NRL/MR/6115--98-8140

Ecosystem-level Evaluation of Intrinsic Contaminant Bioremediation and Impact of Naval Shipyards on Adjacent Ecosystems: A Preliminary Report

M.T. MONTGOMERY

*GEO-Centers, Inc.
Landover, Maryland*

B.J. SPARGO
T.J. BOYD

*Chemical Dynamics and Diagnostics Branch
Chemistry Division*

February 27, 1998

DTIC QUALITY INSPECTED 2

19980305 057

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave Blank)	2. REPORT DATE February 27, 1998	3. REPORT TYPE AND DATES COVERED Interim Report		
4. TITLE AND SUBTITLE Ecosystem-level Evaluation of Intrinsic Contaminant Bioremediation and Impact of Naval Shipyards on Adjacent Ecosystems: A Preliminary Report			5. FUNDING NUMBERS	
6. AUTHOR(S) M.T. Montgomery,* B.J. Spargo and T.J. Boyd				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Research Laboratory Washington, DC 20375-5320			8. PERFORMING ORGANIZATION REPORT NUMBER NRL/MR/6115-98-8140	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) NAVFACENGCOM-South Division, Code 1877 2155 Eagle Drive, North Charleston, SC 29419			10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES *GEO-Centers, Inc., Landover, MD				
12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited.			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) A phase I evaluation of intrinsic contaminant bioremediation was conducted 21-23 July 1997 at a former Defense Reutilization and Marketing Office (DRMO) site at the Charleston Naval Shipyard, Charleston, SC. The objective of the project is to determine the extent to which intrinsic bioremediation is occurring in groundwater contaminated with gasoline (benzene, ethylbenzene, toluene, and xylenes; BTEX), methyl-tertbutyl ether (MTBE), and trichloroethylene (TCE). Preliminary findings suggest that the comingled contaminant plumes have significant bacterial productivity and contaminant mineralization rates. Rapid bacterial production and benzene turnover rates at wells, which are down gradient of the other BTEX plume(s) suggests that the natural bacterial assemblage has the capacity to inhibit migration of the plume towards Noisette Creek. Low ratios of DCE:TCE were found near the expected source but greatly increased as the plume moved towards creek suggesting that <i>in situ</i> biodegradation is occurring. Benzene concentrations did not correlate with changes in bacterial production in the TCE-impacted wells though even in low concentrations. Data suggest there may be three separate BTEX plumes, two from unleaded gasoline, and one from leaded gasoline.				
14. SUBJECT TERMS Bioremediation Groundwater Bacterial productivity Contaminant mineralization			15. NUMBER OF PAGES 34	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT UL	

ECOSYSTEM-LEVEL EVALUATION OF INTRINSIC CONTAMINANT
Bioremediation and Impact of Naval Shipyards on
Adjacent Ecosystem: A Preliminary Report

Michael T. Montgomery, Barry J. Spargo, and Thomas J. Boyd
Environmental Qualities Sciences, Chemistry Division
U.S. Naval Research Laboratory

Abstract

A phase I evaluation of intrinsic contaminant bioremediation was conducted 21-23 July 1997 at a former Defense Reutilization and Marketing Office (DRMO) site at the Charleston Naval Shipyard, Charleston, SC. The objective of the project is to determine the extent to which intrinsic bioremediation is occurring in groundwater contaminated with gasoline (benzene, ethylbenzene, toluene, and xylenes; BTEX), methyl-tertbutyl ether (MTBE), and trichloroethylene (TCE). Preliminary findings suggest that the comingled contaminant plumes have significant bacterial productivity and contaminant mineralization rates. Rapid bacterial production and benzene turnover rates at wells, which are down gradient of the other BTEX plume(s) suggests that the natural bacterial assemblage has the capacity to inhibit migration of the plume towards Noisette Creek. Low ratios of DCE:TCE were found near the expected source but greatly increased as the plume moved towards creek suggesting that *in situ* biodegradation is occurring. Benzene concentrations did not correlate with changes in bacterial production in the TCE-impacted wells though even in low concentrations. Data suggest there may be three separate BTEX plumes, two from unleaded gasoline, and one from leaded gasoline.

Manuscript approved January 20, 1998

Executive Summary

BTEX

- Highest concentrations of BTEX observed at NBCA-039-11
- Bacterial degradation (utilization) of benzene and toluene highest at BTEX plume (NBCA-039-001)
- High benzene turn-over rates and high utilization rates suggest that the plume is rapidly being mineralized by heterotrophic bacteria.
- Low bacterial production (metabolism) coupled with other data suggest that BTEX is the major carbon source
- Rapid bacterial production and benzene turnover rates at wells NBCA-039-009 and -010, which are down gradient of the other BTEX plume(s) suggests that the natural bacterial assemblage has the capacity to inhibit migration of the plume towards Noisette Creek

TCE

- The source of the TCE appears to be near wells NBCA-039-012 and 039-003
- Low ratios of DCE:TCE (Table 1) were found near the expected source but greatly increased as the plume moved towards creek (NBCA039-009, 013) suggesting that *in situ* biodegradation is occurring
- Benzene concentrations did not correlate with changes in bacterial production in the TCE-impacted wells though even in low concentrations
- Low concentrations of TCE correlated with a reduction in heterotrophic activity down to a baseline level

MTBE

- Data suggest there may be three separate BTEX plumes, two from unleaded gasoline (NBCA-039-1 and 039-04), and one from leaded gasoline (NBCA-039-11)
- MTBE is associated with the TCE plume (NBCA-039-012).
- MTBE concentration had no correlation with BTEX concentration
- A strong correlation exists between TCE and MTBE concentration

List of Figures

BTEX

Figure 1. Total BTEX concentration ($\mu\text{g L}^{-1}$) profiles are heavily influenced by MW-11 values.

Figure 2. Ethylbenzene concentration ($\mu\text{g L}^{-1}$) is elevated in MW-11 but in MW-1 in contrast with benzene concentration.

Figure 3. *p*-Xylene concentration ($\mu\text{g L}^{-1}$) is elevated in MW-11 as was concentration of ethylbenzene.

Figure 4. *o*-, *m*-Xylenes concentration ($\mu\text{g L}^{-1}$) is elevated in MW-11 as was concentration of ethylbenzene.

Figure 5. Benzene concentration ($\mu\text{g L}^{-1}$) is elevated in MW-11 and -1 but rapidly decreases with distance.

Figure 6. Benzene utilization ($\mu\text{g L}^{-1} \text{ h}^{-1}$) is highest at MW-1.

Figure 7. Toluene utilization ($\mu\text{g L}^{-1} \text{ h}^{-1}$) is rapid at MW-2 and -7.

Figure 8. Benzene turnover times (h^{-1}) were rapid down gradient of the BTEX spill and in the TCE plume as well as at MW-7. High utilization coupled with fast benzene turnover rates strongly suggest that this plume is rapidly mineralized by heterotrophic bacteria.

Figure 9. Bacterial production in groundwater samples was elevated in the BTEX plume near the Hess border as well as near a plume near buildings 1608A and 1608B. Low bacterial production at this well suggests that BTEX is being used as a relatively major carbon source for heterotrophic bacteria. Generally, the efficiency of carbon transfer to cellular material, during cell growth and reproduction, is lower for recalcitrant aromatic contaminants than for indigenous organic compounds. It might be expected, then, that actively mineralizing populations will have lower relative productivity even though contaminant mineralization is high. This may be confirmed in later studies by stable isotope analyses of CO_2 in the vadose zone and groundwater.

Figure 10. In relatively unimpacted wells, production ($\mu\text{g C L}^{-1} \text{ h}^{-1}$) decreased with increasing benzene concentration ($\mu\text{g L}^{-1}$).

Figure 11. In BTEX impacted wells, production ($\mu\text{g C L}^{-1} \text{ h}^{-1}$) increased at high benzene concentration ($\mu\text{g L}^{-1}$).

Figure 12. In BTEX impacted wells, production ($\mu\text{g C L}^{-1} \text{ h}^{-1}$) increased with increasing benzene concentration ($\mu\text{g L}^{-1}$).

Benzene caused bacterial production to decrease in relatively uncontaminated wells (Fig. 10), but production increases rapidly in the highly BTEX-contaminated wells with benzene (Fig. 11) or total BTEX concentration (Fig. 12)¹. This suggests that as the BTEX plume migrates into an area that is unimpacted by TCE or BTEX, there could be an initial decrease in bacterial metabolism and thus total productivity. In this instance, benzene may either be inhibitory to the natural assemblage or may be less efficiently metabolized as an energy source.

Figure 13. In relatively unimpacted wells, there was little relationship between production ($\mu\text{g C L}^{-1} \text{ h}^{-1}$) and BTEX concentration ($\mu\text{g L}^{-1}$).

Figure 14. In TCE impacted wells, production ($\mu\text{g C L}^{-1} \text{ h}^{-1}$) generally decreased with increasing BTEX concentration ($\mu\text{g L}^{-1}$).

Figure 15. In BTEX impacted wells, benzene utilization ($\mu\text{g C L}^{-1} \text{ h}^{-1}$) decreased with increasing bacterial production ($\mu\text{g C L}^{-1} \text{ h}^{-1}$). In wells with high BTEX contamination, benzene mineralization (utilization) is inversely correlated with bacterial production suggesting that benzene is being inefficiently utilized (rather than inhibitory).

TCE

Figure 16. TCE concentration (relative abundance) is elevated in MW-12 with lower concentration down gradient.

Figure 17. DCE concentration (relative abundance) is elevated in MW-12 with lower concentration down gradient.

Figure 18. Toluene turnover times (h^{-1}) were very rapid in the TCE plume. Low ratios of DCE:TCE (Table 1) were found near the expected source but greatly increased as the plume moved towards creek (NBCA039-009, 013). This evidence coupled with the high toluene turnover rates in the plume (Fig. 18) suggests that the TCE is being degraded by the natural bacterial assemblage. The impact on the creek is difficult to determine without measuring TCE mineralization and groundwater transport rates and possibly having monitoring wells closer to the creek.

Figure 19. In TCE impacted wells, production ($\mu\text{g C L}^{-1} \text{ h}^{-1}$) had little relationship with benzene concentration ($\mu\text{g L}^{-1}$). Unlike in the both unimpacted and BTEX-impacted areas, benzene concentrations did not correlate with changes in bacterial production in the

¹ There was not a clear relationship between total BTEX and production at the unimpacted (Fig. 13) and TCE-impacted wells (Fig. 14).

TCE-impacted wells though even in low concentrations, TCE inhibited bacterial production down to a baseline.

Figure 20. In TCE impacted wells, production ($\mu\text{g C L}^{-1} \text{ h}^{-1}$) decreased with TCE concentration (relative abundance). In the TCE-impacted wells, even low concentrations of TCE correlated with a reduction in heterotrophic activity down to a baseline level ($1.2 \times 10^{-1} \mu\text{g C L}^{-1} \text{ h}^{-1}$)

Figure 21. In TCE impacted wells, toluene utilization ($\mu\text{g C L}^{-1} \text{ h}^{-1}$) increased with increasing bacterial production ($\mu\text{g C L}^{-1} \text{ h}^{-1}$). Though it is possible that TCE inhibits bacterial metabolism of some component of the natural assemblage, it is more likely that the decreased production is a result of reduced metabolic efficiency of toluene degradation. In these wells, heterotrophic production correlated well with toluene utilization. TCE is cometabolized along with toluene metabolism though it is not used as an energy source. This would reduce metabolic efficiency of the natural assemblage and contribute to lower production in the presence of TCE.

MTBE

Figure 22. MTBE concentration ($\mu\text{g L}^{-1}$) is elevated in MW-1, -12, & -14 suggesting at least two sources are present.

Figure 23. In BTEX impacted wells, BTEX concentration ($\mu\text{g L}^{-1}$) had little relationship with MTBE concentration ($\mu\text{g L}^{-1}$). MTBE is associated with the TCE plume (NBCA-039-012). MTBE concentration had no correlation with BTEX concentration suggesting that at least one source of the BTEX was not from unleaded fuel (NBCA-039-011).

Figure 24. In TCE impacted wells, TCE concentration ($\mu\text{g L}^{-1}$) positively correlated with MTBE concentration ($\mu\text{g L}^{-1}$). Surprisingly, there was a strong correlation between TCE and MTBE concentration suggesting that they are co-contaminants with a common source that are being diluted by groundwater transport.

Figure 25. In relatively unimpacted wells, there was some correlation between MTBE concentration ($\mu\text{g L}^{-1}$) and bacterial production ($\mu\text{g C L}^{-1} \text{ h}^{-1}$).

Figure 26. When combining data from all wells, there was little correlation between MTBE concentration ($\mu\text{g L}^{-1}$) and bacterial production ($\mu\text{g C L}^{-1} \text{ h}^{-1}$). There was some increase in bacterial production with MTBE concentration in the relatively unimpacted wells (Fig. 25) though this was not seen when all well data was included (Fig. 26).

List of Tables

Table 1. Contaminant concentrations and biological analyses from July 21-23, 1997 sampling of CNY DRMO site.

Environmental Quality Sciences

Ecosystem-level Evaluation of Intrinsic Contaminant Bioremediation and Impact of Naval Shipyards on Adjacent Environments



Phase I DRMO Site 21-23 July 1997

Total BTEX concn.
($\mu\text{g l}^{-1}$) profiles are
heavily influenced by
MW-11 values.




Figure 1.



Ecosystem-level Evaluation of Intrinsic Contaminant Bio-mediated ation and Impact of Naval Shipyards on Adjacent Environments

Phase I
DRMO Site
21-23 July 1997
Ethylbenzene concn.
($\mu\text{g l}^{-1}$) is elevated in
MW-11 but in MW-1
in contrast with
benzene concn.




Figure 2.



Phase I DRMO Site 21-23 July 1997

p-Xylene concn.
($\mu\text{g l}^{-1}$) is elevated in
MW-11 as was concn.
of ethylbenzene.




Figure 3.

Ecosystem-level Evaluation of Intrinsic Contaminant Bioremediation and Impact of Naval Shipyards on Adjacent Environments

Phase I DRMO Site 21-23 July 1997

o-, *m*-Xylenes concn.
($\mu\text{g l}^{-1}$) is elevated in
MW-11 as was concn.
of ethylbenzene




Figure 4.



Phase I
DRMO Site
21-23 July 1997

Benzene concn. ($\mu\text{g l}^{-1}$)
is elevated in MW-11 &
-1 but rapidly decreases
with distance. Plumes
may not be distinct as
MW-2, -5 were not
sampled.




Figure 5.



Phase I DRMO Site 21-23 July 1997

Benzene utilization
($\mu\text{g l}^{-1} \text{h}^{-1}$) is highest
at MW-1.




Figure 6.

Environmental Quality Sciences

Ecosystem-level Evaluation of Intrinsic Contaminant Bioremediation and Impact of Naval Shipyards on Adjacent Environments



Phase I DRMO Site 21-23 July 1997

Toluene turnover times (h^{-1}) were very rapid in the TCE plume.




Figure 7.



Naval Research Laboratory

Environmental
Quality
Sciences

Ecosystem-level Evaluation of Intrinsic Contaminant Bioremediation and Impact of Naval Shipyards on Adjacent Environments

Phase I DRMO Site 21-23 July 1997

Benzene turnover times (h^{-1}) were rapid downgradient of the BTEX spill and in the TCE plume as well as at MW-7.




Figure 8.



Phase I DRMO Site 21-23 July 1997

Bacterial production (cells $\text{ml}^{-1} \text{h}^{-1}$) in groundwater samples was elevated in the BTEX plume near the Hess border, as well as near a plume by buildings 1608A and 1608B.




Figure 9.




Figure 10.




Figure 11.




Figure 12.




Figure 13.




Figure 14.




Figure 15.

Environmental Quality Sciences

Ecosystem-level Evaluation of Intrinsic Contaminant Bioremediation and Impact of Naval Shipyards on Adjacent Environments



Phase I DRMO Site 21-23 July 1997

TCE concn. (relative abundance) is elevated in MW-12 with lower concn. downgradient.




Figure 16.



Phase I DRMO Site 21-23 July 1997

DCE concn. (relative abundance) is elevated in MW-12 with lower concn. downgradient.




Figure 17.



Ecosystem-level Evaluation of Intrinsic Contaminant Bioremediation and Impact of Naval Shipyards on Adjacent Environments

Phase I
DRMO Site
21-23 July 1997

Toluene turnover
times (h^{-1}) were very
rapid in the TCE
plume.




Figure 18.




Figure 19.




Figure 20.




Figure 21.



Phase I DRMO Site 21-23 July 1997

MTBE concn. ($\mu\text{g l}^{-1}$) is elevated in MW-1, -12, & -14 suggesting at least two sources are present.




Figure 22.




Figure 23.




Figure 24.




Figure 25.




Figure 26.

NBCA 039-	MTBE µg L-1	Benzene µg L-1	Toluene µg L-1	Ethylbenzene µg L-1	o, m-xylene µg L-1	p-xylene µg L-1	Total BTEX µg L-1	DCE µg L-1	TCE µg L-1	DCE:TCE Production	Benzene Utilization µg C L-1 h-1	Toluene Utilization µg C L-1 h-1
1	20.10	69.00	1.05	44.92	59.10	3.13	177.20	0.00	0.00	0.00	1.94E+01	7.33E-01
3	3.26	0.27	0.00	0.49	0.64	0.00	1.40	3.88	30.63	0.13	0.00E+00	1.47E-03
6	0.10	0.45	0.00	0.57	0.00	0.54	1.55	0.00	0.00	0.00	3.30E-01	1.52E-02
7	0.08	0.13	0.00	0.49	0.00	0.00	0.62	0.00	0.00	0.00	7.78E-02	2.61E-01
8	0.00	0.30	0.00	0.80	0.67	0.55	2.32	0.00	0.00	0.00	0.00E+00	9.86E-03
9	0.19	0.33	0.00	0.49	0.00	0.00	0.82	29.46	1.62	18.16	6.84E-02	1.59E-02
10	0.35	0.04	0.00	0.50	0.00	0.62	1.16	0.00	0.00	0.00	0.00E+00	7.11E-03
11	0.00	126.00	0.00	227.00	513.50	54.00	920.50	0.00	0.00	0.00	0.00E+00	0.00E+00
12	28.28	0.29	0.00	0.62	0.67	0.54	2.11	149.02	113.81	1.31	1.15E-01	1.88E-03
13	0.00	0.08	0.00	0.71	0.67	0.00	1.46	23.62	4.51	5.24	6.78E-02	0.00E+00
14	18.63	3.78	0.00	2.78	0.00	3.00	9.55	0.00	0.00	0.00	1.91E+00	0.00E+00
15	0.10	0.13	0.00	0.61	0.00	0.53	1.27	0.00	0.00	0.00	2.52E-01	3.42E-02

Relatively unimpacted wells
Wells highly impacted with TCE
Wells highly impacted with BTEX

Table 1. Contaminant concentrations and biological analyses from July 21-23, 1997 sampling of CNY DRMO site.